



98.5% of patients were satisfied with their vision after the iLASIK® procedure.

Wake up and see without the need for contacts or glasses. With the **iLASIK®** procedure, you can see clearly—any time, anywhere.*

Ask your doctor if it's right for you.

What is the **iLASIK**® procedure?

iLASIK® is a personalized, precise and proven treatment that reshapes the cornea to improve your vision.

iLASIK® is:

- An all-laser, no-blade procedure
- Considered a safe and effective option to reduce or eliminate dependency on glasses or contact lenses
- The iLASIK treatment is quick - takes a few minutes to complete



Personalized.

iLASIK® creates a 3D map of your eye, allowing for a truly customized treatment plan.

Precise.

Majority of eyes achieved 20/16 or better visual acuity.

Proven.

Doctor's #1 choice with over 20 million procedures.[†]

Ask your doctor about the iLASIK® procedure today.



Based on share and shipment data for PIs and Treatment Cards.

Ask your doctor about the **iLASIK®** procedure today.

References:

 Schallhorn SC, Brown M, Venter JA, Teenan D, Hettinger K, Yamamoto H. Early clinical outcomes of wavefront-guided myopic LASIK treatments using a new-generation Hartmann-Shack aberrometer. J Refract Surg. 2014;30(1):14-21.

IMPORTANT SAFETY INFORMATION

INDICATION: LASIK (laser-assisted in situ keratomileusis) is a laser surgery procedure that permanently changes the shape of the cornea to reduce or eliminate nearsightedness, farsightedness, or mixed visual irregularities due to an abnormal curve in the cornea (astigmatism). Only an eye care professional trained in laser vision correction can determine if you are a candidate for this procedure. The iFS® laser is a surgical laser that can be used to create flaps for use in LASIK surgery. CONTRAINDICATIONS: You should not have LASIK if you have collagen vascular (such as rheumatoid arthritis), autoimmune, or an immunodeficiency disease because they affect the body's ability to heal. You should not have this procedure if you are pregnant or nursing, show signs of corneal thinning, or take medications with eye-related side effects, such as Isotretinoin (Accutane®) for acne treatment or Amiodarone hydrochloride (Cordarone®) for normalizing heart rhythm. WARNINGS: LASIK is not recommended if you have diabetes, a history of herpes simplex or herpes zoster keratitis, significant dry eye, or severe allergies. PRECAUTIONS: Your doctor will examine your eyes to determine if you are a candidate for this procedure. Talk to your doctor about any eye-related conditions, injuries, or surgeries you have had, as well as any changes to your vision in the past year. These may result in poor vision after LASIK. Tell your doctor about any medications you are taking. After surgery, you may find it more difficult to see in conditions such as dim light, rain, snow, fog, or glare from bright lights at night. LASIK is for patients 21 years of age and over. SIDE EFFECTS: between 2 and 6 weeks after surgery. Some patients (0.03%) noticed a temporary spoke-like band of light in their peripheral vision. Please consult with your eye care professional and carefully review the Patient Information Booklet regarding the potential risks and benefits of this procedure. Results may vary for each individual patient. The iLASIK® platform utilizes the STAR S4 IR® Excimer Laser System, WaveScan WaveFront® System, as well as the iFS® Advanced Femtosecond Laser during the LASIK procedure. CAUTION: U.S. Federal Law restricts these devices to use by practitioners who have been trained in their calibration and operation, and who have experience in the surgical treatment and management of refractive errors.







Johnson Johnson vision